CONTENTS

NQR studies on 2,5-dichlorophenol
A. Kasturi, P. Venkatacharyulu and D. Premaswarup (Nagaranjunagar, India) 1
Long-range substituent effects on conformational equilibria
O. Exner (Prague, Czechoslovakia), M.J. Cook and T.J. Howe (Norwich, Gt. Britain) 11
Hydrogen-bonding effect on ^{15}N NMR chemical shifts of the glycine residue of oligopeptides in the solid state as studied by high-resolution solid-state NMR spectroscopy
S. Kuroki, S. Ando and I. Ando (Tokyo, Japan), A. Shoji and T. Ozaki (Gunma, Japan) and G.A. Webb (Guildford, Gt. Britain) 19
Chirality in *trans*-1,4-cyclohexane dicarboxylic acid
M. Barón (Buenos Aires, Argentina) 31
Phase transition in solid 2,2,4,4-tetramethylpentan-3-ol (TMP)
G. Bator, R. Jakubas, Z. Malarski, Z. Galewski, J. Matuszewski and A. Miniewicz (Wrocław, Poland) 39
An unusual geometrical and spectroscopic isotopic effect in the solid complex of 4-methylpyridine with pentachlorophenol
I. Majerz, Z. Malarski and T. Lis (Wrocław, Poland) 47
β-Diketone interactions. Part 12. The structure and properties of 3-(3′,4′,5′-trimethylphenyl)-pentane-2,4-dione, $C_{14}H_{18}O_2$; the hydrogen-bond energy of the enol tautomers of β-diketones
J. Emsley, L.Y.Y. Ma, S.C. Nyburg and A.W. Parkins (London, Gt. Britain) 59
Crystal structure of bis(betaine)hydrochloride monohydrate
X.-M. Chen and T.C.W. Mak (New Territories, Hong Kong) 69
Crystal and molecular structures of the first *trans*-MACRO-ANSA monocyclophosphazene macrocycle, *trans*-$N_3P_3Cl_4[HN–(CH_2)_3–O–(CH_2)_6–O–(CH_2)_3–NH]$
B. Zanin, F. Sournies, J.-F. Labarre, R. Enjalbert and J. Galy (Toulouse, France) 77
Tetraosmium carbonyl clusters: an X-ray photoelectron spectroscopy (XPS) study of valence and core levels
R. Zanoni (Rome, Italy) and J. Puga (Midland, MI, U.S.A.) 89
The molecular structure of 1,2,3-trichlorobenzene determined by combined analysis of data from gas electron diffraction and liquid crystal NMR and by X-ray crystallography
A.J. Blake, R. Blom, S. Cradock, S.L. Davidson and D.W.H. Rankin (Edinburgh, Gt. Britain) 95
Crystallographic studies on sterically affected chemical species. Part II. Molecular and crystal structure of 1,8-bis(dimethylamino)naphthalene tetrafluoroborate. Analysis of distortion of geometry in the aromatic part due to intramolecular hydrogen bonding
K. Woźniak and T.M. Krygowski (Warsaw, Poland), B. Kariuki and W. Jones (Cambridge, Gt. Britain) and E. Grech (Szczecin, Poland) 111
Crystallographic studies of intra- and inter-molecular interactions. Part IV. Molecular and crystal structure of phthalazine semi-tetrafluoroborate: intramolecular changes in geometry as a consequence of H-bonding
K. Woźniak and T.M. Krygowski (Warsaw, Poland), B. Kariuki and W. Jones (Cambridge, Gt. Britain) and E. Grech (Szczecin, Poland) 119
Study of γ-irradiated chloral hydrate single crystal by electron spin resonance
R. Krzyminiewski and J. Kasprzak (Poznań, Poland) 127
An ESR study of the stable radical in a γ-irradiated single crystal of 17α-hydroxy-progesterone
R. Krzyminiewski, J. Pietrzak and R. Konopka (Poznań, Poland) 133
Molecular structure and conformation of diisopropyl sulfide in the vapor phase: reinvestigation by molecular orbital constrained electron diffraction
H. Takeuchi, S. Konaka (Sapporo, Japan), K. Siam (Pittsburg, KS, U.S.A.), L. Schäfer and J.D. Ewbank (Fayetteville, AR, U.S.A.) 141

Distances, root-mean-square amplitudes of vibration, and vibrational anharmonicity in carbon tetrabromide from a gas-phase electron-diffraction investigation
H. Thomassen and K. Hedberg (Corvallis, OR, U.S.A.) 151
A quantitative comparison of spectroscopic and energy data, in solution, of NHO and OHO hydrogen bonds and gas-phase proton affinities: complexes of pyridines and pyridine *N*-oxides with acetic acids
Z. Dega-Szafran, A. Hrynio and M. Szafran (Poznań, Poland) 159
IR spectroscopic investigation of solvent effects and H-bond formation on the CD stretching frequency of chloroform
H. Kleeberg, C. Eisenberg and T. Zinn (Marburg, F.R.G.) 175
Fourier transform-infrared spectra and molecular orientation in thin Langmuir–Blodgett films of dodecyloxyphenylpyrazine carboxylic acid and its barium salt
T. Kamata, J. Umemura, T. Takenaka (Kyoto, Japan), K. Takehara, K. Isomura and H. Taniguchi (Fukuoka, Japan) 187
Conformational change of diethylene glycol diethyl ether due to cation-capture and water-accommodation
K. Fukushima and K. Sakurada (Shizuoka, Japan) 201
Fourier transform-infrared study of hydrogen-bonding between an OH-donor and a vicinal oxygen acceptor
F.A.J. Singelenberg and J.H. van der Maas (Utrecht, The Netherlands) 213
Conformational stability, structure and vibrational spectra of cyclopropylmethanol
H.M. Badawi, M.E. Abu-Zeid and Y.A. Yousef (Safat, Kuwait) 225
Infrared matrix isolation studies of dimethylsulphoxide–hydrogen iodide molecular complexes
A.J. Barnes (Salford, Gt. Britain), L. Schriver, A. Schriver and J.-P. Perchard (Paris, France) 239
Microwave spectrum and amino hydrogen location in indole
W. Caminati and S. Di Bernardo (Bologna, Italy) 253
Investigation of the molecular structure of catechol by combined microwave spectroscopy and ab initio calculations
W. Caminati, S. Di Bernardo (Bologna, Italy), L. Schäfer and S.Q. Kulp-Newton (Fayetteville, AR, U.S.A.) and K. Siam (Pittsburg, KA, U.S.A.) 263
Lattice vibration spectra. Part LVIII. OD(OH) frequency versus H-bond distance correlation diagrams: a case study for isomorphic $M(SO_3)_2 \cdot 3H_2O$ (M = Mg, Mn, Fe, Co, Ni, Zn)
H.D. Lutz, J. Henning and B. Engelen (Siegen, F.R.G.) 275
Monomeric and dimeric copper(II) compounds with the 2-fluoro-6-chlorobenzoate ligand
M. Melnik, Z. Sabova (Bratislava, Czechoslovakia), J. Mroziński and A. Szymaszek (Wrocław, Poland) 285
Model potential functions of nonlinear XY_2 molecules
K.V. Ermakov, B.S. Butayev and V.P. Spiridonov (Moscow, U.S.S.R.) 295
Orbital interactions in diaziridines
S.N. Denisenko, G. Kaupp (Oldenburg, F.R.G.), A.J. Bittner and P. Rademacher (Essen, F.R.G.) 305
Conformational analyses of the isomers of the $[Co\{(\pm)pn\}_n(en)_{3-n}]^{3+}$ systems: a reinvestigation (en = ethane-1,2-diamine; pn = propane-1,2-diamine; n = 0–3)
I.M. Atkinson, F.R. Keene (Townsville, Qld., Australia) and G.H. Searle (Adelaide, S.A., Australia) 313
Solution-state conformation of 1,1′-binaphthyl. A Kerr and Cotton–Mouton effect study
P. Wormell and G.L.D. Ritchie (Armidale, N.S.W., Australia) 331

Book reviews 339

Author index 343

Subject index 344